

## INFESTATION OF TAILLESS AMPHIBIANS OF GENUS RANA BY TREMATODES IN THE VALLEY OF THE TISA RIVER (YUGOSLAVIA)

ESTER POPOVIĆ and M. MIKEŠ

Institute of Biology, Novi Sad, Yugoslavia  
(Received September 11, 1988)

### Abstract

On a section comprising the territory of Bečej and Ečka a parasitological search has been carried out in 13 hosts of the species *Rana esculenta* and 9 individuals of the species *Rana ridibunda*. Almost the same extensity of invadedness in both species of frogs has been stated (77.8% and 76.9%), while the specimens of the species *Rana ridibunda* have shown a greater intensivity of infestation. Nine species of trematodes have been defined: *Diplodiscus subclavatus* GOETE, *Gorgodera cygnoides* ZEDER, *Haematoloechus (Pneumonoeces) variegatus* RUDOLPHI, *Haematoloechus (Pneumonoeces) schulzei* WUNSCH, *Opisthyoglyphe ranae* FRÖLICH, *Cephalogonimus retusus* DUJARDIN, *Pleurogenoides medians* OLSSON, *Prostotocus confusus* LOOSS and *Pleurogenes claviger* RUDOLPHI.




Fig. 1 — *Diplodiscus subclavatus*  
Fig. 2 — *Gorgodera cygnoides*  
Fig. 3 — *Opisthyoglyphe ranae*



Fig. 4 — *Cephalogonimus retusus*

Fig. 5 — *Pleurogenes claviger*

Fig. 6 — *Pleurogenoides medians*




Fig. 7 — *Prosotocus confusus*

## Introduction

Presently the interest for the investigation of parasitofauna has been growing constantly both with respect to their place in the systems and their spread, as well as from the ecological aspect. The basic problem in the domain of ecology is, of course, to state how the parasites develop in the interaction on the line exterior environment-parasite-host, as well as to define their role in given ecosystems, in this case, primarily in the aquatic ones. Namely due to alimentary concatenation, amphibians, particularly the hosts of endohelminths, are organically connected to a great deal with the members of marshy biocenoses, among others with many economically important species.

The investigation of helminthofauna of tetrapodes from the open air of Vojvodina (the northern part of Yugoslavia) so far has been sporadically carried out, and that of the hosts of ornitofauna (SEY *et al.* 1971; SOTI *et al.* 1972; MIKES *et al.* 1974), of terriofauna — the mouselike rodents from agrobiocenoses (MÉSZÁROS *et al.* 1983), from small mammals of the periodically inundated zone of the Tisa River (MIKES, HABIJAN, MIKES, 1986), as well as *Anura*, also from the valley of the Tisa River (MIKES, POPOVIĆ 1988). Data on helminthofauna Amphibia in Yugoslavia can be found only in works HRISTOVSKI 1968, 1974 from the territory of Macedonia, and trematodes of the small cormorant (*Phalacrocorax pygmaeus*) from the territory of the Scutari Lake have been treated by POPOVIĆ (in press).

In this paper we present the infestation of two species of hosts — the tailless amphibians from genus *Rana* (*R. esculenta*)-13 frogs and *R. ridibunda*-9 frogs by trematodes. Beside defining the level of the quantitative and qualitative invadence of the hosts by certain groups of helminths, the analysis of the collected material is also concerned with the defining of the taxonomic belonging of the discovered trematodes.

### Materials and Methods

The analysis of the infestation of the hosts by endohelminths has been performed on a total of 22 individuals of tailless amphibians (13 ind. *Rana esculenta* and 9 ind. *R. ridibunda*). The hosts originate from two specific localities in the valley of the Tisa River. One locality is situated in the valley of the Tisa River near Bečej. The biotope itself represents a littoral zone of the river, which is overgrown by *Amorpha fruticosa* in the frame of *Saliceto-Populetum nigrae*, a component which is situated in a narrow girdle of only 30—150 m between the protective dam and the river. The other locality comprises the periodically inundated territory of the estuary of the Begej into the Tisa River with Carska Bara and several large anthropogenous ponds. The analysed material originates from the pit of Carska bara surrounded by *Saliceto-Populetum nigrae* components, in front of which there is a very developed *Scripo-Phragmitetum* marshy community and a floatant association *Nymphaeetum albo-luteae*.

After having noted the data on localities, the data of catch, and elaborated the host (taxonomic belonging, biometric data, sex and age composition), we have submitted each individual to the parasitological examination by standard method. The search of the host with respect to the infestation by endoparasitic helminths has been performed at the level of lungs, urinary bladder and at some levels of the digestive tract (Gaster, intestinum, rectum).

The collected parasites have been elaborated macrotechnically, and conserved in 70% alcohol, in order to be elaborated and defined microtechnically later on. Permanent preparations have been stained with alau-carmin and fixed by Canada balsam. The defining has been performed according to the EDELENYI key (1974).

### The extensity and intensivity of the infestation by indigenous trematodes

The results of the analysis of infestation in general show a high degree of infestation by trematodes (about 3/4 of the examined individuals are invaded), and that almost with the same extensity in both species of hosts (Tab. 1). When analysing the

distribution of parasites according to their localization, it has been stated that trematodes occur mainly in intestinum and lungs, and that in the species *R. ridibunda* in a somewhat higher percentage. The extensity of infestation shows a high level of infected condition in both species of hosts — in the case of *R. esculenta* there is in the intestinum a four times greater number, and in *R. ridibunda* a two times greater number of trematodes in relation to the infestation of lungs.

The infestation of the host by trematodes with respect to the intensivity of the invadedness of organs shows a relatively low level of the infestation of lungs in relation to intestinum (Table 1). It may be noticed that the high level of invadedness

Tab. 1. The locality of invasion of the host with Trematods

Host	№			Organum													
				Intestinum						Pulmo				Ves. urin.			
	EXP	INF	%	№	%	1—10	11—20	36—45	51—60	71—75		%	1—5	16—10	№	%	1—5
<i>Rana esculenta</i>	13	10	76,9	8	80,0	2	3	2	1	2	20,0	2		1	10,0	1	
<i>Rana ridibunda</i>	9	7	77,8	6	85,7	1	2		2	1	3	42,8	2	1			

of the intestinum makes its appearance simultaneously with an important intensivity in the category of above 50 percent trematodes per host, and that with a somewhat increased number of individuals in the species *R. ridibunda*.

Eight species of trematodes, out of nine defined species, have been stated in the host *R. esculenta*, and in *R. ridibunda*. It has to be added that liver-flukes *Haematolechus schulzei* is absent in the species *R. esculenta*, and the liver-fluke *Gorgodera cygnoides* has not been stated in the host *R. ridibunda* (Table 2). The remaining seven

Tab. 2. The extensity and intensity of the invasion of the host with Trematods

Trematodes	Rana esculenta										Rana ridibunda									
	Extensity					Intensity					Extensity					Intensity				
	№					№					№					№				
	EXP	INF	%	1-5	6-10	11-15	31-35	51-55			EXP	INF	%	1-5	6-10	11-15	16-20	21-25	46-51	
1. <i>Diplodiscus subclavatus</i>	10	3	30,0	2	1						7	6	85,7	6						
2. <i>Gorgodera cygnoides</i>	10	2	20,0	2							7	—								
3. <i>Haemotolechus variegatus</i>	10	1	10,0	1							7	2	28,6	2						
4. <i>Haematolechus schulzei</i>	10										7	2	28,6	2						
5. <i>Opisthyoglyphe ranae</i>	10	7	70,0	5	1		1				7	3	42,8			1	1		1	
6. <i>Clephalogonimus retusus</i>	10	6	60,0	2	1	1	1	1			7	6	85,7	3	1			1	1	
7. <i>Pleurogenes claviger</i>	10	1	10,0	1							7	2	28,6	2						
8. <i>Pleurogenoides medians</i>	10	1	10,0	1							7	1	14,3	1						
9. <i>Prosotocus confusus</i>	10	1	10,0	1							7	2	28,6		2					

species of trematodes have been found in both examined species of hosts. The liver-flukes and *Opisthyoglyphe ranae* and *Cephalogonimus retusus* in both species of hosts show a high level of the extensity of infestation (from 42.8% to 85.7%), and the species *Diplodiscus subclavatus* is present in the case of *R. ridibunda*.

### Faunistic and taxonomic survey of defined species of trematodes

In the two examined species of Anura, 9 species of trematodes (belonging to 5 families) have been stated. It makes the half of the already known 18 species of trematodes in frogs. The following species of trematodes have been determined:

1. *Diplodiscus subclavatus* (PALLAS, 1790), GOEZE, 1782

Host: *Rana esculenta* L. — discovered in 3 individuals

*Rana ridibunda* PALLAS — discovered in 6 individuals

Locality: Bečej

Localization: intestinum

2. *Gorgodera cygnoides* ZEDER

Host: *Rana esculenta* L. — 2 infested individuals

Locality: Ečka, Bečej

Localization: vesica urinaria

3. *Haematoloechus variegatus* RUDOLPHI

Host: *Rana esculenta* L. — 1 infested individual

*Rana ridibunda* PALLAS — 2 infested individuals

Locality: Bečej

Localization: pulmo

4. *Haematoloechus schulzei* WUNDSCH

Host: *Rana ridibunda* PALLAS — 2 infested individuals

Locality: Bečej

Localization: pulmo

5. *Optisthyoglyphe ranae* FRÖHLICH

Host: *Rana esculenta* L. — 7 infested hosts

*Rana ridibunda* PALLAS — 3 infested hosts

Locality: Bečej

Localization: intestinum

6. *Cephalogonimus retusus* DUJARDIN

Host: *Rana esculenta* L. — 6 infested hosts

*Rana ridibunda* PALLAS — 6 infested hosts

Locality: Bečej

Localization: intestinum

7. *Pleurogenes claviger* RUDOLPHI

Host: *Rana esculenta* L. — infested individual

*Rana ridibunda* PALLAS — 2 infested individuals

Locality: Bečej

Localization: intestinum



8. *Pleurogenoides medians* OLSSON  
Host: *Rana esculenta* L. — 1 infested individual  
*Rana ridibunda* PALLAS — 1 infested individual  
Locality: Bečej  
Localization: intestine
9. *Prostocus confusus* LOOSS  
Host: *Rana esculenta* L. — 1 infested individual  
*Rana ridibunda* PALLAS — 2 infested individuals  
Locality: Bečej  
Localization: intestine

### Morphological features of the defined trematodes

#### 1. *Diplodiscus subclavatus*

The length of the body is 1.4 mm—3.5 mm, and the width 0.58—1.27 mm. The body is pear-shaped. The abdominal sucker is terminal, very developed. In the middle part of the abdominal sucker, another sucker may be seen. The testis is a little greater than the oral sucker  $316.0\text{--}647.8 \times 276.5\text{--}474.0\text{ }\mu\text{m}$ . The ovary is smaller than the testis  $110.6\text{--}260.7 \times 118.5\text{--}244.9\text{ }\mu\text{m}$ . The dimensions of the eggs  $86.9\text{--}158 \times 63.2\text{--}86.9\text{ }\mu\text{m}$ .

#### 2. *Gorgodera cygnoides*

The length of the body is 1.8 mm and 5.79 mm, and the width 0.49 mm, 1.22 mm). Oral sucker has smaller dimensions ( $221.2\text{ }\mu\text{m}$  and  $434.5\text{ }\mu\text{m} \times 244.9\text{ }\mu\text{m}$  and  $505.6\text{ }\mu\text{m}$ ) than the abdominal one ( $363.4\text{ }\mu\text{m}$  and  $948.0\text{ }\mu\text{m} \times 379.2\text{ }\mu\text{m}$  and  $897\text{ }\mu\text{m}$ ). Testis are not clearly divided in the left and right group ( $126.4\text{--}395.0\text{ }\mu\text{m} \times 189.6\text{--}537.2\text{ }\mu\text{m}$ ). The dimensions of the ovary are  $173.8\text{ }\mu\text{m}$  and  $521.4\text{ }\mu\text{m} \times 237.0\text{ }\mu\text{m}$  and  $442.4\text{ }\mu\text{m}$ . The eggs are oval:  $30.0\text{--}36.25 \times 18.75\text{--}25.0\text{ }\mu\text{m}$ .

#### 3. *Haematoloechus variegatus*

The length of the body is 5.0—9.0 mm, and the width 0.916—1.659 mm. The surface of the body does not possess any pricks. Oral sucker is subterminal ( $355.5\text{--}655.7 \times 260.7\text{--}647.8\text{ }\mu\text{m}$ ). The abdominal sucker has smaller dimensions ( $244.9\text{--}308.1 \times 252.8\text{--}308.1\text{ }\mu\text{m}$ ). The testis are elongated-oval. They are situated on the medial line, one behind the other ( $T_1 = 553.0\text{--}813.7 \times 450.3\text{--}521.4\text{ }\mu\text{m}$ ;  $T_2 = 529.3\text{--}1.343 \times 347.6\text{--}584.6\text{ }\mu\text{m}$ ). The dimensions of the ovary:  $323.9\text{--}884.8 \times 268.5\text{--}790.0\text{ }\mu\text{m}$ , and those of the eggs:  $30.0\text{--}51.25 \times 15.0\text{--}30.0\text{ }\mu\text{m}$ .

#### 4. *Haematoloechus Schulzei*

The length of the body is 3.1—5.5 mm, and the width 1.098—1.437 mm. The surface of the body is covered with tiny pricks. The dimensions of the oral sucker:  $292.3\text{--}418.7 \times 371.3\text{--}434.5\text{ }\mu\text{m}$ , and the abdominal sucker  $308.1\text{--}323.9 \times 308.1\text{--}355.5\text{ }\mu\text{m}$ . The first testis  $553.0\text{--}639.9 \times 371.3\text{--}521\text{ }\mu\text{m}$ , and the second testis:  $711.0\text{--}726.8 \times 276.5\text{--}521.4\text{ }\mu\text{m}$ . The ovary ( $379.2\text{--}474.0 \times 237.0\text{--}395.0\text{ }\mu\text{m}$ ) is situated directly below the abdominal sucker. The dimensions of eggs:  $30.0\text{--}52.5 \times 15.0\text{--}30.0\text{ }\mu\text{m}$ .

#### 5. *Opisthyoglyphe ranae*

The dimensions of the body are  $0.853-1.532 \times 0.331-0.726$  mm. The surface of the body is covered by small pricks. Oral sucker is larger ( $110.6-181.7 \times 118.5-189.0 \mu\text{m}$ ) than the abdominal one ( $92.5-165.9 \times 96.25-134.3 \mu\text{m}$ ). Testis are situated in the posterior half of the body medially one behind the other ( $T_1 = 79.0-158.0 \times 115.0-284.4 \mu\text{m}$ ;  $T_2 = 79.0-165.9 \times 108.75-276.5 \mu\text{m}$ ). Ovary ( $75.0-197.5 \times 71.25-173.8 \mu\text{m}$ ) is situated at left side of the body, near the abdominal sucker. The dimensions of the eggs:  $36.25-56.25 \times 22.5-35.0 \mu\text{m}$ .

#### 6. *Cephalogonimus retusus*

The length of the liver-fluke is  $1.319-3.500$  mm, and the width  $0.395-0.853$  mm. Tiny pricks cover the surface of the body. Oral sucker ( $181.7-268.6 \times 181.7-284.4 \mu\text{m}$ ) is larger than abdominal one ( $122.5-205.4 \times 133.75-229.7 \mu\text{m}$ ). The dimensions of the first testis:  $101.25-237.00 \times 112.5-308.1 \mu\text{m}$ , and the second testis:  $110.6-268.6 \times 125.0-252.8 \mu\text{m}$ . Ovary is situated at the left side of the body ( $76.25-229.1 \times 66.25-181.7 \mu\text{m}$ ). The eggs:  $31.25-65.0 \times 15.0-27.5 \mu\text{m}$ .

#### 7. *Pleurogenes claviger*

The length of the body is  $1.830$  mm and  $2.725$  mm, and the width  $0.774$  mm,  $0.963$  mm. Pricks cover the surface of the body. The dimensions of the oral sucker:  $205.4 \times 237.0 \mu\text{m}$  and  $237.0 \times 316.0 \mu\text{m}$ , and the abdominal one:  $150.1 \times 158.0 \mu\text{m}$ , and  $189.6 \times 237.0 \mu\text{m}$ . The size of the first testis:  $244.9 \times 276.5 \mu\text{m}$ , and  $316.0 \times 244.9 \mu\text{m}$ , and the second:  $213.3 \times 237.0 \mu\text{m}$ ;  $252.8 \times 284.4 \mu\text{m}$ . The length of the ovary:  $237.0 \mu\text{m}$  and the width  $252.8 \mu\text{m}-268.6 \mu\text{m}$ . The dimensions of the eggs:  $27.5-31.25 \times 75-17.5 \mu\text{m}$ .

#### 8. *Pleurogenoides medians*

The length of the body:  $0.671$  mm, and the width  $0.363$  mm. The dimensions have been registered only in one trematode. The dimensions of oral sucker:  $95.0 \times 110.0 \mu\text{m}$ , and the abdominal one:  $97.5 \times 96.25 \mu\text{m}$ , and  $118.5 \times 126.4 \mu\text{m}$ . The body is covered with tiny pricks. The length, and the width of the first testis:  $134.3 \times 165.9 \mu\text{m}$  and  $158.0 \times 110.6 \mu\text{m}$ , and the second testis:  $126.4 \times 110.6 \mu\text{m}$  and  $173.8 \times 158.0 \mu\text{m}$ . The dimensions of the ovary:  $107.5 \times 80.0 \mu\text{m}$ , and  $158.0 \times 134.3 \mu\text{m}$ . The eggs:  $15.0-31.25 \times 11.25-16.25 \mu\text{m}$ .

#### 9. *Prosotocus confusus*

The length of the body:  $0.813-1.350$  mm, and the width  $0.513-0.861$  mm. Tiny pricks cover the surface of the body. Oral sucker is larger ( $142.2-237.0 \times 165.9-221.2 \times 112.5-237.0 \mu\text{m}$ ). The dimension of the right testis:  $110.6-237.0 \times 158.0-237.0 \mu\text{m}$ ; that of the left one:  $134.3-221.2 \times 158.0-252.8 \mu\text{m}$ . The length of the ovary:  $134.3-221.2 \mu\text{m}$  and the width  $158.0-237.0 \mu\text{m}$ . The dimensions of the eggs:  $17.5-30.0 \times 10.0-15.0 \mu\text{m}$ .

### Conclusion

The parasitological examination of two species of hosts of the genus *Rana* (*R. esculenta* and *R. ridibunda*) from the valley of the river Tisa, with respect to the infestation by trematodes, has given the following results:

— The presence of nine species of parasites, belonging to five families of trematodes has been stated, what makes the half of the already known species of trematodes in Anura. Two of them

— *Haematoloechus schulzei* is absent in the host *R. esculenta*, and the trematode *Gorgoderina cygnoides* has not been stated in the species *R. ridibunda* (Table 2).

— First of all, trematodes appear in the intestinum and lungs of the host. As to their distribution in hosts, in the intestinum of *R. esculenta* a four times greater number of trematodes has been stated, while in the lungs of *R. ridibunda* a two times greater number of trematodes has been found.

— Simultaneously with the high level of infestation there appears a significant intensivity in the category above 50 percent trematodes per host, and that with a somewhat higher number of individuals in the species *R. ridibunda* (Tab. 2).

— The extensity of invadedness of hosts by trematodes is significant. In both species of hosts about 3/4 of examined individuals are infested (Table 1). A high degree of invadedness with *Opisthyoglyphe ranae* (42.8%) and *Cephalogonimus retusus* has been stated in both species of hosts and with *Diplodiscus subclavatus* (85.7%) in the case of the species *R. ridibunda* (Tab. 2).

## References

- BRELIH, S., DŽUKIĆ, G. (1974): Catalogus faunae Jugoslaviae — Reptilia IV/2. — SAZU, Ljubljana.
- DELY, O. (1967): Kétféltűek — Amphibia, Fauna Hungariae 83. — Akad. kiadó, Budapest.
- EDELÉNYI, B. (1974): Mételyek II — Trematodes II. Közzetett fejlődésű mételyek — Digenea, — Akad. kiadó, Budapest.
- HRISTOVSKI, N. (1968): *Gorgoderina alobata* LEES i Mitohell 1966, nova vrsta (Trematoda — Gorgoderidae) za faunata na Jugoslavia (*Gorgoderina alobata* LEES et Mitschell, 1966 a sp. nov. (Trematoda — Gorgoderidae) to the fauna of Yugoslavia), — Prilozi, 9: 45—48, Bitola.
- HRISTOVSKI, N. (1979): The nematodes of the suborder Oxyurata in the vicinity of Bitola, — Pril. 30—31, 135—157, Bitola.
- MÉSZÁROS, F., HABIJAN, V., MIKES, M. (1983): Parasitic nematodes of rodents in Vojvodina (Yugoslavia), — Parasit. Hung. 16, 103—110.
- MIKES, M., SOTI, J., SEY, O., DIMITRIJEVIĆ, S. (1974): Survey of the trematodes of the fish-eating birds in Vojvodina, — Arh. biol. 26, 55—59.
- MIKES, M., HABIJAN, V., MIKES, B. (1986): Prikaz infestiranosti teriofaune endohelminthima u dolini reke Tise (The infestedness of teriofauna by endohelminths in the valley of the Tisa river). — Zb. za prir. n. MS, 71, 145—154.
- MIKES, M., POPOVIĆ, E. (1988): Infestiranost nekih vrsta *Anura helminthima* duž reke Tisa (The infestation of some species of *Anura* by *helminths* along the river Tisa), — Zb. rad. PMF, ser. biol. 18, 35—47.
- POPOVIĆ, E.: Mali kormoran (*Phalacrocorax pygmaeus* PALL., 1773) domaćin Echinostomatidnih metilja (Trematodes: Echinostomatidae) (Pygmy Cormorant — *Phalacrocorax pygmaeus* PALL., 1773 as a host of the Echinostomatid trematodes — Trematodes: Echinostomatidae) — (in press).
- RADOVANOVIĆ, M. (1952): Vodozemci i gmizavci naše zemlje (Amphibians and Reptiles of Yugoslavia), — NK Beograd.
- SEY, O., MIKES, M., SOTI, J., DIMITRIJEVIĆ, S. (1974): Trematodes of birds occurred in Vojvodina, — Zb. rad. PMF I, 193—210.
- SOTI, J., SEY, O., MIKES, M. (1972): Trematodák és Cestodák vajdasági madarakból, — Acta Acad. Pedag. in Civit. Pécs 16, 13—24.



## A Tiszavölgy Rana (Amphibia) békáinak Trematoda fertőzöttsége (Yugoslavia)

POPOVIĆ ESZTER és MIKES M.

Egyetemi Biológiai Intézet, Újvidék

### Kivonat

A Tiszavölgy Becse és Écska térségéről gyűjtött *Rana* nemhez tartozó 13 *Rana esculenta* és 9 *Rana ridibunda* helmintológiai vizsgálatát végezték el a szerzők. Megállapítást nyert, hogy a gazdaállatok egyedeinek fertőzöttségi extenzitása, mindkét békafaj esetében, többé-kevésbé egyenletes (77,8% és 76,9%). Ugyanakkor a tavi béka fertőzöttségi intenzitása elenyészően nagyobb. Összesen 9 Trematoda faj került elő: *Diplodiscus subclavatus* GOETE, *Haematoloechus (Pneumonoeces) variegatus* RUDOLPHI, *Haematoloechus (Pneumonoeces) schulzei* WUNDSCH, *Opisthyoglyphe ranae* FRÖLICH, *Gorgodera cygnoides* ZEDER, *Cephalogonimus retusus* DUJARDIN, *Pleurogenes claviger* RUDOLPHI, *Pleurogenoides medians* OLSSON és *Prosotocus confusus* LOOSS.

## Инфестированность безхвостых земноводных рода (Amphibia: Anura) в долине реки Тиса

Эстер Попович и М. Микеш

Институт Биологии Естественного-математического Факультета г. Нови Сад

### Резюме

На участке реки Тиса, в районе г. Бечей и м. Эчка, выполнены паразитологические испытания 13 экземпляров породы *Rana esculenta* и 9 экземпляров породы *Rana ridibunda*. Установлена была почти идентичная экстенсивность инвадированности подвергаемых анализу примеров обоих пород лягушек (77,8%, т.е. 76,9%), в то время как у экземпляров пород *R. ridibunda* обнаружена увеличенная интенсивность зараженности. Установлено было 9 пород Trematoda: *Diplodiscus subclavatus* GOETE, *Gorgodera cygnoides* ZEDER, *Haematoloechus (Pneumonoeces) variegatus* RUDOLPHI, *Haematoloechus (P.) schulzei* WUNDSCH, *Opisthyoglyphe ranae* FRÖLICH, *Cephalogonimus retusus* DUJARDIN, *Pleurogenoides medians* OLSSON, *Prosotocus confusus* LOOSS и *Pleurogenes claviger* RUDOLPHI.

## Infestiranost bezrepih vodozemaca roda Rana (Amphibia: Anura) u dolini reke Tise

ESTER POPOVIĆ i M. MIKEŠ

Institut za biologiju, Novi Sad

### Izvod

Na deonici reke Tise, sa područja Bečeja i Ečke, izvršena je parazitološka pretraga 13 domaćina vrste *Rana esculenta* i 9 jedinki vrste *Rana ridibunda*. Ustanovljen je skoro istovetan ekstenzitet invadiriranosti kod analiziranih jedinki obe vrste žabe (77,8% odnosno 76,9%), dok su primerci vrste *R. ridibunda* imala veći intenzitet zaraženosti. Determinisano je 9 vrsta Trematoda: *Diplodiscus subclavatus* GOETE, *Gorgodera cygnoides* ZEDER, *Haematoloechus (Pneumonoeces) variegatus* RUDOLPHI, *Haematoloechus (P.) schulzei* WUNDSCH, *Opisthyoglyphe ranae* FRÖLICH, *Cephalogonimus retusus* DUJARDIN, *Pleurogenoides medians* OLSSON i *Prosotocus confusus* LOOSS, i *Pleurogenes claviger* RUDOLPHI.